Leaflet KEBENE ANTIACID: 150x210 mm

K SHTANTZA

KEBENE® antiacid

1,2 g
Vanilla taste

Directions for use

Triple action:
Symptomatic treatment of gastric pain and heartburn
Reduces gastroesophageal reflux
Antibloating effect

Chewable tablets
Vanilla taste
Lactose free
Gluten free

INGREDIENTS
Calcium carbonate, Magnesium hydroxide, Simethicone, Sorbitol, Sodium citrate, Magnesium stearate, Aroma, Hydroxypropylcellulose, Maltodextrin, Silicon dioxide, Sucralose.

This product contains an amount of 855 mg of calcium carbonate, 120 mg of magnesium hydroxide, 120 mg of sodium citrate and 60 mg of simethicone for a single intake dose (equal to 2 tablets of 1.2 grams).

INDICATIONS
KEBENE ANTIACID is indicated in the treatment of burning and gastric pain caused by excess acidity (reduces gastric hyperacidity). The product is also useful in treating symptoms associated with gastroesophageal reflux and dyspepsia (such as trouble digesting, gastric burning and epigastric pain (feeling full right after a meal).

HOW IT WORKS
KEBENE ANTIACID is a medical device in chewable tablets form, containing a combination of calcium carbonate, magnesium hydroxide, sodium citrate and simethicone. Once arrived in the stomach, the product neutralizes the pH of gastric juices and acid vapors, exerting a protective and soothing action on the gastro-esophageal mucosa, with a rapid action against epigastric pain due to hyperacidity. Simethicone is a substance that decreases the surface tension of the gas bubbles in the gastrointestinal tract, facilitating their expulsion, reducing the feeling of bloating and abdominal distension.

HOW TO TAKE AND POSOLOGY
Adults and children over 12 years of age: It is advisable to take 2 tablets as needed (in case you experience burning and / or pain symptoms) to be taken after main meals and in the evening before bedtime. Chew well before swallowing. Do not exceed the maximum dose of 8 tablets per day.

WARNINGS
Adults and children over 12 years of age: It is advisable to take 2 tablets as needed (in case you experience burning and / or pain symptoms) to be taken after main meals and in the evening before bedtime. Chew well before swallowing. Do not exceed the maximum dose of 8 tablets per day.
Please consult the doctor in case of hypercalcaemia (high levels of calcium in the blood), hypermagnesemia (high levels of magnesium in the blood), nephrocalcinosis and recurrent calcium containing renal calculi.
There are no contraindications to use during pregnancy and lactation, but it is advisable in this case to seek medical advice before use. Do not use if the package is open or damaged, or in the presence of visible changes in the appearance of the tablets.
Do not use the product after the expiry date printed on the package.

SIDE EFFECTS
After taking the product mild and transient side effects may occur, such as: bloating, abdominal pain, belching, flatulence, constipation.
The use of the product in compliance with the instructions reduces the risk of unwanted side effects.

Leaflet KEBENE ANTIACID: 150x210 mm

K SHTANTZA

HOW TO STORE
Keep out of the reach and sight of children. The product must be stored at a temperature between 8 ° C and 25 ° C, in a cool and dry place and away from heat sources. The expiry date refers to the product in intact packaging and properly stored. Do not store tablet parts for later use

PACK
KEBENE ANTIACID is presented in blister form containing 40 chewable tablets of 1.2 g each.

Biofarma – Via Castelliere 2 – Mereto di Tomba (UD) – Italia

Distributed by: Terapia SA
Str. Fabricii nr. 124, Cluj-Napoca, Romania

Medical Device CE 0373

Date of last revision: August 2023

(IFU1_08/23r0_08/23)